

PRIMES IN THE FORM $\lfloor \alpha p + \beta \rfloor$

HONGZE LI AND HAO PAN

ABSTRACT. Let β be a real number. Then for almost all irrational $\alpha > 0$ (in the sense of Lebesgue measure)

$$\limsup_{x \rightarrow \infty} \pi_{\alpha, \beta}^*(x) (\log x)^2 / x \geq 1,$$

where

$$\pi_{\alpha, \beta}^*(x) = \{p \leq x : \text{both } p \text{ and } \lfloor \alpha p + \beta \rfloor \text{ are primes}\}.$$

Recently Jia [4] solved a conjecture of Long and showed that for any irrational number $\alpha > 0$, there exist infinitely many primes not in the form $2n + 2\lfloor \alpha n \rfloor + 1$, where $\lfloor x \rfloor$ denotes the largest integer not exceeding x . Subsequently, in [2] Banks and Shparlinski investigated the distribution of primes in the Beatty sequence $\{\lfloor \alpha n + \beta \rfloor : n \geq 1\}$. Motivated by the binary Goldbach conjecture and the twin primes conjecture, we have the following conjecture:

Conjecture 1. *Let $\alpha > 0$ be an irrational number and β be a real number. Then there exist infinitely many primes p such that $\lfloor \alpha p + \beta \rfloor$ is also prime.*

On the other hand, Deshouillers [3] proved that for almost all (in the sense of Lebesgue measure) $\gamma > 1$ there exist infinitely many primes p in the form $\lfloor n^\gamma \rfloor$. Furthermore, Balog [1] showed that for almost all $\gamma > 1$

$$\limsup_{x \rightarrow \infty} \frac{|\{p \leq x : \text{both } p \text{ and } \lfloor p^\gamma \rfloor \text{ are primes}\}|}{x/(\log x)^2} \geq \gamma.$$

In this note we shall show that Conjecture 1 holds for almost all α . Define

$$\pi_{\alpha, \beta}^*(x) = \{p \leq x : \text{both } p \text{ and } \lfloor \alpha p + \beta \rfloor \text{ are primes}\}.$$

Theorem 1. *Let β be a real number. Then*

$$\limsup_{x \rightarrow \infty} \pi_{\alpha, \beta}^*(x) (\log x)^2 / x \geq 1 \tag{1}$$

for almost all irrational $\alpha > 0$.

For a set $X \subseteq \mathbb{R}$, let $\text{mes}(X)$ denote its Lebesgue measure. Without the additional mentions, the constants implied by $\ll$, $\gg$ and $O(\cdot)$ will be always absolute.

2000 *Mathematics Subject Classification.* Primary 11N05; Secondary 11N36, 11P32.

This work was supported by the National Natural Science Foundation of China (Grant No. 10771135).

Lemma 1. *Let $I \subseteq [0, 1)$ be an interval. Suppose that $b, l > 0$. Then*

$$\text{mes}(\{\alpha \in (0, b) : \{\alpha/l\} \in I\}) = O(b+l)\text{mes}(I),$$

where $\{x\} = x - \lfloor x \rfloor$.

Proof. Without loss of generality, we may assume that $I = (c_1, c_2)$ with $0 \leq c_1 < c_2 \leq 1$. Let $J = \{\alpha \in (0, b) : \{\alpha/l\} \in I\}$. Clearly

$$J \subseteq \bigcup_{0 \leq j \leq b/l} ((j + c_1)l, \min\{b, (j + c_2)l\}).$$

If $l \leq b$, then

$$\text{mes}(J) \leq (\lfloor b/l \rfloor + 1)(c_2 - c_1)l = O(b)\text{mes}(I).$$

And if $l > b$, then

$$\text{mes}(J) = \text{mes}((c_1l, \min\{b, c_2l\})) \leq (c_2 - c_1)l.$$

□

Lemma 2. *Suppose that $b_2 > b_1 > 0$ and β are arbitrarily real numbers. Let $\epsilon > 0$ be a small number and x be a sufficiently large (depending on b_1, b_2, β and ϵ) integer. Then there exists an exceptional set $J_E \subseteq (b_1, b_2)$ with $\text{mes}(J_E) = O(x^{-\epsilon})$ such that for any square-free $d \leq x^{1/3-2\epsilon}$ and irrational $\alpha \in (b_1, b_2) \setminus J_E$,*

$$|\{1 \leq n \leq x : n\lfloor \alpha n + \beta \rfloor \equiv 0 \pmod{d}\}| = \frac{x}{d} \prod_{p|d} \left(2 - \frac{1}{p}\right) + O(x^{1-\epsilon}/d). \quad (2)$$

Proof. For an irrational $\alpha \in (b_1, b_2)$, let

$$\mathcal{A}(x; \alpha) = \{n\lfloor \alpha n + \beta \rfloor : 1 \leq n \leq x\}$$

and

$$\mathcal{A}_d(x; \alpha) = \{a \in \mathcal{A} : a \equiv 0 \pmod{d}\}.$$

For a square-free d , we have

$$\begin{aligned} |\mathcal{A}_d(x; \alpha)| &= \sum_{s|d} |\{1 \leq n \leq x/s : \lfloor \alpha sn + \beta \rfloor \equiv 0 \pmod{d/s}, (n, d/s) = 1\}| \\ &= \sum_{s|d} \sum_{t|d/s} \mu(t) |\{1 \leq n \leq x/s : \lfloor \alpha sn + \beta \rfloor \equiv 0 \pmod{d/s}, t | n\}| \\ &= \sum_{s|d, t|s} \mu(t) |\{1 \leq n \leq x/s : \lfloor \alpha sn + \beta \rfloor \equiv 0 \pmod{dt/s}\}|. \end{aligned}$$

Clearly

$$\lfloor \alpha sn + \beta \rfloor \equiv 0 \pmod{dt/s} \iff \{\alpha ns^2/td + \beta s/td\} \in [0, s/td).$$

Let $\alpha' = \alpha s^2/td$, $\beta' = \beta s/td$, $d' = td/s$ and $y = x/s$. Clearly $y \geq x^{2/3+2\epsilon}$ and $d' \leq d$. Let

$$I_{a,q} = \{\theta \in [0, 1) : |\theta q - a| \leq x^{2\epsilon}/y\}.$$

Suppose that $d'x^{2\epsilon} \leq q \leq y/x^{2\epsilon}$ and $1 \leq a \leq q$ with $(a, q) = 1$. If $\{\alpha'\} \in I_{a,q}$,

$$\begin{aligned} & |\{1 \leq n \leq y : \{an/q + \beta'\} \in [1/q, 1/d' - 1/q]\}| \\ & \leq |\{1 \leq n \leq y : \{\alpha'n + \beta'\} \in [0, 1/d']\}| \\ & \leq |\{1 \leq n \leq y : \{an/q + \beta'\} \in [0, 1/d' + 1/q] \cup [1 - 1/q, 1]\}|. \end{aligned}$$

Hence

$$|\{1 \leq n \leq y : \{\alpha'n + \beta'\} \in [0, 1/d']\}| = y/d' + O(q/d') + O(y/q).$$

Let

$$\mathcal{I}_{d'} = \bigcup_{\substack{1 \leq a \leq d'x^{2\epsilon} \\ (a,q)=1}} I_{a,q}$$

Clearly

$$\text{mes}(\mathcal{I}_{d'}) \leq \sum_{\substack{1 \leq a \leq d'x^{2\epsilon} \\ (a,q)=1}} \text{mes}(I_{a,q}) \ll \frac{d'x^{4\epsilon}}{y} = tdx^{4\epsilon-1}.$$

If $\alpha s^2/td \notin \mathcal{I}_{td/s}$ for each s, t with $s \mid d, t \mid s$, then

$$|\mathcal{A}_d(x; \alpha)| = \sum_{\substack{s \mid d, t \mid s}} \mu(t)x/td(1 + O(x^{-2\epsilon})) = \frac{x}{d} \prod_{p \mid d} (2 - 1/p) + O(x^{1-\epsilon}/d).$$

Let

$$\mathcal{J}_d = \{\alpha \in (0, b) : \{\alpha s^2/td\} \in \mathcal{I}_{td/s} \text{ for some } s, t \text{ with } s \mid d, t \mid s\}.$$

Applying Lemma 1,

$$\text{mes}(\mathcal{J}_d) \ll b_2 \sum_{\substack{s \mid d, t \mid s \\ b_2 \geq td/s^2}} \text{mes}(\mathcal{I}_{td/s}) + \frac{td}{s^2} \sum_{\substack{s \mid d, t \mid s \\ b_2 < td/s^2}} \text{mes}(\mathcal{I}_{td/s}) = O(x^{-1/3+\epsilon}).$$

Finally, Let

$$J_E = \bigcup_{d \leq x^{1/3-2\epsilon}} \mathcal{J}_d.$$

Clearly we have $\text{mes}(J_E) = O(x^{-\epsilon})$. □

Lemma 3. Suppose that $b_2 > b_1 > 0$, $\epsilon > 0$ and β are arbitrarily real numbers. Then there exists an exceptional set $J_E \subseteq (b_1, b_2)$ with $\text{mes}(J_E) = O(x^{-\epsilon})$ such that for any irrational $\alpha \in (b_1, b_2) \setminus J_E$,

$$|\{1 \leq p \leq x : \text{both } p \text{ and } \lfloor \alpha p + \beta \rfloor \text{ are primes}\}| \ll \frac{x}{(\log x)^2} \quad (3)$$

for sufficiently large (depending on b_1, b_2, β and ϵ) x .

Proof. Let $z = x^{1/8}$. Define

$$P(z) = \prod_{\substack{p < z \\ p \text{ prime}}} p$$

and

$$\mathcal{S}(A, z) = \{a \in A; (a, P(z)) = 1\}.$$

Let $\mathcal{A}(\alpha) = \{n \lfloor \alpha n + \beta \rfloor : 1 \leq n \leq x\}$. Clearly

$$\{p \lfloor \alpha p + \beta \rfloor : z + \alpha^{-1}(z + 1 - \beta) \leq p \leq x, \text{ both } p \text{ and } \lfloor \alpha p + \beta \rfloor \text{ are primes}\}$$

is a subset of $\mathcal{S}(\mathcal{A}(\alpha), z)$. Furthermore, by Lemma 2, we know that there exists a set $J_E \subseteq (b_1, b_2)$ with $\text{mes}(J_E) = O(x^{-\epsilon})$ such that for any square-free $1 \leq d \leq x^{1/3-2\epsilon}$ and irrational $\alpha \in (b_1, b_2) \setminus J_E$,

$$|\mathcal{A}_d(\alpha)| = \frac{x}{d} \prod_{p|d} \left(2 - \frac{1}{p}\right) + O(x^{1-\epsilon}/d),$$

where $\mathcal{A}_d(\alpha) = \{y \in \mathcal{A}(\alpha) : d \mid y\}$. Let $g(m)$ be the completely multiplicative function such that $g(p) = 2/p - 1/p^2$ for each prime p . Define $G(z) = \sum_{\substack{m < z \\ m|P(z)}} g(m)$.

By Selberg's sieve method,

$$|\mathcal{S}(\mathcal{A}(\alpha), z)| \leq \frac{|\mathcal{A}(\alpha)|}{G(z)} + O\left(\sum_{\substack{d < z^2 \\ d \text{ square-free}}} 3^{\omega(d)} x^{1-\epsilon}/d\right),$$

where $\omega(d)$ denotes the number of distinct prime divisors of d . Since $3^{\omega(d)} \ll d^\epsilon$,

$$\sum_{d < z^2} \frac{3^{\omega(d)}}{d} \ll z^{2\epsilon}.$$

So it suffices to show $G(z) \gg (\log z)^2$. By Theorem 7.14 in [5], we know

$$G(z) = \sum_{\substack{m < z \\ m|P(z)}} g(m) \gg \prod_{p < z} (1 - g(p))^{-1} = \prod_{p < z} (1 - 2/p + 1/p^2)^{-1} \gg (\log z)^2.$$

□

Proof of Theorem 1. Suppose that $b_2 > b_1 > 0$. Let

$$\mathcal{F} = \{\alpha \in (b_1, b_2) : \limsup_{x \rightarrow \infty} \pi_{\alpha, \beta}^*(x)(\log x)^2/x < 1\}$$

and

$$\mathcal{F}_n = \{\alpha \in (b_1, b_2) : \limsup_{x \rightarrow \infty} \pi_{\alpha, \beta}^*(x)(\log x)^2/x \leq 1 - 1/n\}.$$

Clearly $\mathcal{F} = \bigcup_{n > 1} \mathcal{F}_n$. So it suffices to show that $\text{mes}(\mathcal{F}_n) = 0$ for every $n > 1$. (The measurability of $\mathcal{F}_n$ will be proven later.)

Assume on the contrary that there exists $n > 1$ such that $\text{mes}(\mathcal{F}_n) > 0$. Let $I = (c_1, c_2)$ be an arbitrary sub-interval of (b_1, b_2) . Clearly

$$\begin{aligned}
\int_{c_1}^{c_2} \pi_{\alpha, \beta}^*(x) d\alpha &= \int_{c_1}^{c_2} \left(\sum_{\substack{p \leq x \\ p \text{ prime}}} \sum_{\substack{\alpha p + \beta - 1 < q \leq \alpha p + \beta \\ q \text{ prime}}} 1 \right) d\alpha \\
&= \sum_{\substack{p \leq x \\ p \text{ prime}}} \sum_{\substack{c_1 p + \beta - 1 < q \leq c_2 p + \beta \\ q \text{ prime}}} \text{mes}([(q - \beta)/p, (q + 1 - \beta)/p] \cap [c_1, c_2]) \\
&\geq \sum_{\substack{p \leq x \\ p \text{ prime}}} \sum_{\substack{c_1 p + \beta < q \leq c_2 p + \beta - 1 \\ q \text{ prime}}} \frac{1}{p} \\
&= (c_2 - c_1) \sum_{\substack{p \leq x \\ p \text{ prime}}} \frac{1}{\log p} \left(1 + O\left(\frac{1}{\log(c_1 p)}\right) \right) \\
&\geq (c_2 - c_1) \frac{x}{(\log x)^2} \left(1 + O\left(\frac{1}{\log x}\right) \right), \tag{4}
\end{aligned}$$

provided that x is sufficiently large (depending on b_1 and b_2). Suppose that $C > 1$ is the implied constant in Lemma 3. Let $\mathcal{L}_I = \mathcal{F}_n \cap I$ and

$$\mathcal{L}_{I, \delta}(x) = \{\alpha \in I : \pi_{\alpha, \beta}^*(x) \leq (1 - \delta)x/(\log x)^2\}.$$

For any two primes p and q , clearly

$$J_{p, q} := \{\alpha \in I : \lfloor \alpha p + \beta \rfloor = q\}$$

is an interval or empty set. Hence

$$\mathcal{L}_{I, \delta}(x) = I \setminus \left(\bigcup_{\substack{k > (1 - \delta)x/(\log x)^2 \\ p_1, \dots, p_k \leq x \text{ are distinct primes} \\ q_1, \dots, q_k \text{ are primes}}} \bigcap_{j=1}^k J_{p_j, q_j} \right)$$

is measurable in the sense of Lebesgue measure. Let $\epsilon > 0$ be a very small number. By Lemma 3,

$$\begin{aligned}
\int_{c_1}^{c_2} \pi_{\alpha, \beta}^*(x) d\alpha &\leq O(x^{1 - \epsilon}) + \\
&+ \text{mes}(\mathcal{L}_{I, \delta}(x)) \frac{(1 - \delta)x}{(\log x)^2} + (c_2 - c_1 - \text{mes}(\mathcal{L}_{I, \delta}(x))) \frac{Cx}{(\log x)^2} \tag{5}
\end{aligned}$$

provided that x is sufficiently large. Combining (4) and (5), we have

$$\text{mes}(\mathcal{L}_{I, \delta}(x)) \leq \frac{C - 1}{C - 1 + \delta/2} \text{mes}(I). \tag{6}$$

We claim that

$$\mathcal{L}_I = \bigcap_{m>n} \bigcup_{y \geq 1} \bigcap_{x \geq y} \mathcal{L}_{I, 1/n-1/m}(x). \quad (7)$$

In fact, for any $m > n$, if

$$\limsup_{x \rightarrow \infty} \frac{\pi_{\alpha, \beta}^*(x)}{x/(\log x)^2} < 1 - \frac{1}{n} + \frac{1}{m},$$

then there exists y_0 such that for any $x \geq y_0$

$$\pi_{\alpha, \beta}^*(x) \leq \left(1 - \frac{1}{n} + \frac{1}{m}\right) \frac{x}{(\log x)^2}.$$

On the other hand, if $\alpha \in \bigcup_y \bigcap_{x \geq y} \mathcal{L}_{I, 1/n-1/m}(x)$, clearly we have

$$\limsup_{x \rightarrow \infty} \frac{\pi_{\alpha, \beta}^*(x)}{x/(\log x)^2} \leq 1 - \frac{1}{n} + \frac{1}{m}.$$

By (6) and (7), we get

$$\text{mes}(\mathcal{L}_I) \leq \limsup_{x \rightarrow \infty} \mathcal{L}_{I, 2/3n}(x) \leq \frac{C-1}{C-1+1/3n} \text{mes}(I).$$

Since $\text{mes}(\mathcal{F}_n) > 0$, there exist open intervals $I_1, I_2, \dots \subseteq (b_1, b_2)$ such that

$$\mathcal{F}_n \subseteq \bigcup_{k=1}^{\infty} I_k$$

and

$$\sum_{k=1}^{\infty} \text{mes}(I_k) \leq \frac{C-1+1/4n}{C-1} \text{mes}(\mathcal{F}_n).$$

But by (6),

$$\text{mes}(\mathcal{F}_n) = \sum_{k=1}^{\infty} \text{mes}(\mathcal{L}_{I_k}) \leq \frac{C-1}{C-1+1/n} \sum_{k=1}^{\infty} \text{mes}(I_k) \leq \frac{C-1+1/4n}{C-1+1/3n} \text{mes}(\mathcal{F}_n).$$

This evidently leads to a contradiction. $\square$

Remark. In [6] and [8], Harman proved that for almost all real $\alpha > 0$ there are infinitely many pairs of (p, q) satisfying

$$|\alpha p - q| < \psi(p), \quad p, q \text{ are primes,}$$

provided that ψ is a non-increasing positive function and

$$\sum_{\substack{2 \leq p \leq \infty \\ p \text{ primes}}} \frac{\psi(p)}{\log p} \quad (8)$$

diverges. (In fact, in [8] Harman established a quantitative version of the above result, on condition that $\psi(n) \in (0, 1/2)$ for each n .) As an immediate consequence,

for almost all $\alpha > 0$, there exists infinitely many pair of primes (p, q) such that $\llbracket \alpha p \rrbracket = q$, where $\llbracket x \rrbracket$ is the nearest integer to x . For more related results, the readers may refer to [7, Chapter 6].

Acknowledgment. We are grateful to Professor Glyn Harman for his very helpful discussions and kindly sending us the copies of the references [6] and [8].

REFERENCES

- [1] A. Balog, *On a variant of the Piatetski-Shapiro prime number problem*. Publ. Math. Orsay, 1989, 3-11.
- [2] W. D. Banks and I. E. Shparlinski, *Prime numbers with Beatty sequences*, preprint, arXiv:0708.1015.
- [3] J. Deshouillers, *Nombres premiers de la forme $[n^c]$* . C. R. Acad. Sci., Paris, Ser. A, **282** (1976), 131-133.
- [4] C.-H. Jia, *On a conjecture of Yiming Long*. Acta Arith., **122** (2006), 57-61.
- [5] C.-D. Pan and C.-B. Pan, *Goldbach Conjecture*. Science Press, Beijing, 1992.
- [6] G. Harman, *Metric diophantine approximation with two restricted variables. III: Two prime numbers*. J. Number Theory, **29** (1988), 364-375.
- [7] G. Harman, *Metric number theory*. London Mathematical Society Monographs. New Series 18. Oxford, Clarendon Press. 1998.
- [8] G. Harman, *Variants of the second Borel-Cantelli lemma and their applications in metric number theory*. Bambah, R. P. (ed.) et al., Number theory. Basel, Birkhäuser. Trends in Mathematics. 121-140, 2000.

E-mail address: lih@sjtu.edu.cn

E-mail address: haopan79@yahoo.com.cn

DEPARTMENT OF MATHEMATICS, SHANGHAI JIAOTONG UNIVERSITY, SHANGHAI 200240,
PEOPLE'S REPUBLIC OF CHINA